Exercise Set 5.2

- 1. Use mathematical induction (and the proof of Proposition 5.2.1 as a model) to show that any amount of money of at least 14¢ can be made up using 3¢ and 8¢ coins.
- 2. Use mathematical induction to show that any postage of at least 12¢ can be obtained using 3¢ and 7¢ stamps.
- 3. For each positive integer n, let P(n) be the formula

$$1^{2} + 2^{2} + \dots + n^{2} = \frac{n(n+1)(2n+1)}{6}$$

- a. Write P(1). Is P(1) true?
- b. Write P(k).
- c. Write P(k + 1).
- d. In a proof by mathematical induction that the formula holds for all integers $n \ge 1$, what must be shown in the inductive step?
- 4. For each integer *n* with $n \ge 2$, let P(n) be the formula

$$\sum_{i=1}^{n-1} i(i+1) = \frac{n(n-1)(n+1)}{3}.$$

- a. Write P(2). Is P(2) true?
- b. Write P(k).
- c. Write P(k+1).
- d. In a proof by mathematical induction that the formula holds for all integers $n \ge 2$, what must be shown in the inductive step?
- 5. Fill in the missing pieces in the following proof that

$$1 + 3 + 5 + \dots + (2n - 1) = n^2$$

for all integers $n \ge 1$.

Proof: Let the property P(n) be the equation

$$1+3+5+\cdots+(2n-1)=n^2. \leftarrow P(n)$$

Show that P(1) is true: To establish P(1), we must show that when 1 is substituted in place of *n*, the left-hand side equals the right-hand side. But when n = 1, the left-hand side is the sum of all the odd integers from 1 to $2 \cdot 1 - 1$, which is the sum of the odd integers from 1 to 1, which is just 1. The right-hand side is (a), which also equals 1. So P(1) is true.

Show that for all integers $k \ge 1$, if P(k) is true then P(k+1) is true: Let k be any integer with $k \ge 1$.

[Suppose P(k) is true. That is:]

Suppose $1 + 3 + 5 + \dots + (2k - 1) = \underline{(b)}$. $\leftarrow P(k)$ [This is the inductive hypothesis.]

[We must show that P(k + 1) is true. That is:]

We must show that

1

$$\underline{(c)} = \underline{(d)}, \qquad \leftarrow P(k+1)$$

But the left-hand side of P(k + 1) is

$$+3+5+\dots+(2(k+1)-1) = 1+3+5+\dots+(2k+1)$$
 by algebra
= $[1+3+5+\dots+(2k-1)] + (2k+1)$
the next-to-last term is $2k-1$ because (e)
= $k^2 + (2k+1)$ by (f)
= $(k+1)^2$ by algebra

which is the right-hand side of P(k + 1) [as was to be shown.]

[Since we have proved the basis step and the inductive step, we conclude that the given statement is true.]

The previous proof was annotated to help make its logical flow more obvious. In standard mathematical writing, such annotation is omitted.

Prove each statement in 6-9 using mathematical induction. Do not derive them from Theorem 5.2.2 or Theorem 5.2.3.

- 6. For all integers $n \ge 1$, $2+4+6+\cdots+2n = n^2+n$.
- 7. For all integers $n \ge 1$,

$$1+6+11+16+\dots+(5n-4)=\frac{n(5n-3)}{2}.$$

- 8. For all integers $n \ge 0$, $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} 1$.
- 9. For all integers $n \ge 3$,

$$4^3 + 4^4 + 4^5 + \dots + 4^n = \frac{4(4^n - 16)}{3}.$$

Prove each of the statements in 10-17 by mathematical induction.

- 10. $1^2 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$, for all integers $n \ge 1$.
- 11. $1^3 + 2^3 + \dots + n^3 = \left[\frac{n(n+1)}{2}\right]^2$, for all integers $n \ge 1$.
- 12. $\frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \dots + \frac{1}{n(n+1)} = \frac{n}{n+1}$, for all integers $n \ge 1$.
- 13. $\sum_{i=1}^{n-1} i(i+1) = \frac{n(n-1)(n+1)}{3}$, for all integers $n \ge 2$.

14.
$$\sum_{i=1}^{n+1} i \cdot 2^{i} = n \cdot 2^{n+2} + 2$$
, for all integers $n \ge 0$.

H 15.
$$\sum_{i=1}^{n} i(i!) = (n+1)! - 1$$
, for all integers $n \ge 1$.

- 16. $\left(1-\frac{1}{2^2}\right)\left(1-\frac{1}{3^2}\right)\cdots\left(1-\frac{1}{n^2}\right)=\frac{n+1}{2n}$, for all integers $n \ge 2$.
- 17. $\prod_{i=0}^{n} \left(\frac{1}{2i+1} \cdot \frac{1}{2i+2} \right) = \frac{1}{(2n+2)!}, \text{ for all integers } n \ge 0.$
- **4 * 18.** If x is a real number not divisible by π , then for all integers $n \ge 1$,

 $\sin x + \sin 3x + \sin 5x + \dots + \sin (2n-1)x$

$$=\frac{1-\cos 2nx}{2\sin x}$$

19. (For students who have studied calculus) Use mathematical induction, the product rule from calculus, and the facts that $\frac{d(x)}{dx} = 1$ and that $x^{k+1} = x \cdot x^k$ to prove that for all integers $n \ge 1$, $\frac{d(x^n)}{dx} = nx^{n-1}$.

Use the formula for the sum of the first n integers and/or the formula for the sum of a geometric sequence to evaluate the sums in 20–29 or to write them in closed form.

20.
$$4 + 8 + 12 + 16 + \dots + 200$$

21. $5 + 10 + 15 + 20 + \dots + 300$

22.
$$3 + 4 + 5 + 6 + \dots + 1000$$

- 23. $7 + 8 + 9 + 10 + \dots + 600$
- **24.** $1 + 2 + 3 + \dots + (k 1)$, where k is an integer and $k \ge 2$.
- **25.** a. $1 + 2 + 2^2 + \dots + 2^{25}$ b. $2 + 2^2 + 2^3 + \dots + 2^{26}$
- 26. $3 + 3^2 + 3^3 + \cdots + 3^n$, where *n* is an integer with $n \ge 1$
- 27. $5^3 + 5^4 + 5^5 + \dots + 5^k$, where *k* is any integer with $k \ge 3$.

28.
$$1 + \frac{1}{2} + \frac{1}{2^2} + \dots + \frac{1}{2^n}$$
, where *n* is a positive integer

- 29. $1 2 + 2^2 2^3 + \dots + (-1)^n 2^n$, where *n* is a positive integer
- **H** 30. Find a formula in n, a, m, and d for the sum $(a + md) + (a + (m + 1)d) + (a + (m + 2)d) + \dots + (a + (m + n)d)$, where m and n are integers, $n \ge 0$, and a and d are real numbers. Justify your answer.
 - 31. Find a formula in a, r, m, and n for the sum

$$ar^m + ar^{m+1} + ar^{m+2} + \cdots + ar^{m+n}$$

where *m* and *n* are integers, $n \ge 0$, and *a* and *r* are real numbers. Justify your answer.

- 32. You have two parents, four grandparents, eight greatgrandparents, and so forth.
 - a. If all your ancestors were distinct, what would be the total number of your ancestors for the past 40 generations (counting your parents' generation as number one)? (*Hint:* Use the formula for the sum of a geometric sequence.)
 - b. Assuming that each generation represents 25 years, how long is 40 generations?
 - c. The total number of people who have ever lived is approximately 10 billion, which equals 10¹⁰ people. Compare this fact with the answer to part (a). What do you deduce?

258 Chapter 5 Sequences, Mathematical Induction, and Recursion

Find the mistakes in the proof fragments in 33-35.

H 33. Theorem: For any integer $n \ge 1$,

$$1^2 + 2^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

"Proof (by mathematical induction): Certainly the theorem is true for n = 1 because $1^2 = 1$ and $\frac{1(1+1)(2 \cdot 1+1)}{6} = 1$. So the basis step is true. For the inductive step, suppose that for some integer $k \ge 1$, $k^2 = \frac{k(k+1)(2k+1)}{6}$. We must show that $(k+1)^2 = \frac{(k+1)((k+1)+1)(2(k+1)+1)}{6}$."

H 34. Theorem: For any integer $n \ge 0$,

 $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1.$

"Proof (by mathematical induction): Let the property P(n) be $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$. Show that P(0) is true: The left-hand side of P(0) is $1 + 2 + 2^2 + \dots + 2^0 = 1$

and the right-hand side is $2^{0+1} - 1 = 2 - 1 = 1$ also. So P(0) is true."

H 35. Theorem: For any integer $n \ge 1$,

$$\sum_{i=1}^{n} i(i!) = (n+1)! - 1.$$

"Proof (by mathematical induction): Let the property P(n) be $\sum_{i=1}^{n} i(i!) = (n+1)! - 1$.

Show that P(1) is true: When n = 1

So
$$\sum_{i=1}^{1} i(i!) = (1+1)! - 1$$
$$1(1!) = 2! - 1$$
$$1 = 1$$

Thus P(1) is true."

- ★ 36. Use Theorem 5.2.2 to prove that if m and n are any positive integers and m is odd, then ∑_{k=0}^{m-1}(n + k) is divisible by m. Does the conclusion hold if m is even? Justify your answer.
- $H \neq 37$. Use Theorem 5.2.2 and the result of exercise 10 to prove that if p is any prime number with $p \ge 5$, then the sum of squares of any p consecutive integers is divisible by p.